

Autophagy as new emerging cellular effect of nanomaterials

ZHONG WenYing^{1†}, LÜ Min^{1†}, LIU LiYing^{2,3}, SUN JinLi^{2,3}, ZHONG ZengTao³,
ZHAO Yun^{1*} & SONG HaiYun^{2*}

¹ School of Life Science, Sichuan University, Chengdu 610064, China;

² Institute for Nutritional Sciences, Shanghai Institutes for Biological Sciences, Shanghai 200031, China;

³ College of Life Sciences, Nanjing Agricultural University, Nanjing 210095, China

Received March 27, 2013; accepted May 6, 2013

The biosafety issue of nanoscale materials is getting more and more attention with their increasing manufacture and application. In the research of cellular effects and underlying mechanisms related to toxicity of nanomaterials, most emphasis were placed on processes such as apoptosis, metabolic inhibition and oxidative stress. Recent evidence suggests that autophagy is part of the biological effects by nanomaterials and various kinds of nanomaterials are capable of disturbing the autophagic process. This review will highlight the importance of autophagy as an emerging mechanism of nanomaterial toxicity and the implication in the therapy of autophagy-related diseases. We summarize current research status of interaction between nanomaterials and autophagic pathways. It is of note that nanomaterials can either induce or block autophagy, which result in similar phenotype but completely different biological consequence. It is therefore important to perform comprehensive analysis of the whole autophagic flux in the future research.

autophagy, nanomaterials, cytotoxicity, cellular effect

Citation: Zhong W Y, Lü M, Liu L Y, et al. Autophagy as new emerging cellular effect of nanomaterials. *Chin Sci Bull*, 2013, 58: 4031–4038, doi: 10.1007/s11434-013-6058-x

Nanomaterials refer to materials with the size range between 1 and 100 nanometers in one or more dimensions [1]. Due to their superior physicochemical properties, such as ultra small size, increased ratio of surface area to volume, multiple options for modification and relatively good biocompatibilities, application of nanomaterials in research as well as in industry becomes more and more common. It is predicted that the total amount of nanomaterials' production will increase up to 58000 tons by 2020. Particularly, the utilization of nanomaterials in biomedical research as biosensors, drug-carriers, or imaging agents was extensively studied [2–4]. Large scaled preparation and application of nanomaterials increase their possibilities to be exposed to human beings or enter the environment [5–7]. Therefore the impact of nanomaterials on environment and human health has attracted high attention. Related studies were carried out

on the biological effect of various kinds of nanomaterials both *in vitro* and *in vivo* [8]. To date, reported physiological changes of a cell or an organism that exposed to nanomaterials include: reactive oxygen species (ROS)-related pathological responses (such as damaged membrane, mitochondrial damage and necrosis), apoptosis, inflammation and altered differentiation patterns, etc [9–12].

Recently, autophagy as an essential cellular process and a novel biological effect of nanomaterials on eukaryotes, has received much attention. Many nanomaterials were reported to be able to change the basal level of autophagy [13,14]. Autophagy can crosstalk with other cellular process, such as apoptosis and necrosis, and therefore is an important part of the cellular effect of nanomaterials [15,16]. Moreover, nanomaterials-induced autophagy showed potential for the treatment of certain diseases including neurodegenerative diseases and cancer [13,17]. Hence, in this review we summarize the current research on the effect of various kinds of nanomaterials on autophagy. We divide the nanomaterials

†These authors contributed equally to this work.

*Corresponding authors (email: hysong@sibs.ac.cn; zhaoyun@scu.edu.cn)

into several categories according to their chemical composition: (1) Carbon-based nanomaterials, (2) metal-based nanomaterials, (3) semiconductor-based nanoparticles, (4) rare earth containing nanomaterials.

1 Autophagy

The homeostasis of cells requires a balance between synthetic and catabolic processes. As one of the catabolic pathways, autophagy mainly regulated the intracellular degradation of long-lived or mis-folded proteins and damaged organelles [18–20]. In most cells, autophagy occurs constitutively at low level and plays an important role in the cellular metabolism, growth and development [21]. Autophagy can be induced by either physiological stress conditions or exogenous stimuli including chemicals and invading particles.

Recently, various kinds of nanomaterials were reported to be able to induce autophagy [17,22–25]. Based on the different mode of intracellular substrates entering lysosome, there are three forms of autophagy: macroautophagy, microautophagy and chaperone-mediated autophagy. Autophagy induced by nanomaterials often refers to macroautophagy, which was firstly identified by De Duve in 1963 [26]. Macroautophagic process occurs in three stages: (1) The formation of double-membrane-structured vesicles (autophagosome) that containing damaged proteins and organelles. At this stage, microtubule-associated protein light chain 3 (LC3) convert from a cytoplasmic form (LC3-I) to a phosphatidylethanolamine (PE)-conjugated form (LC3-II) and localize onto the membrane of autophagosomes. This process is often triggered by drugs (for example, rapamycin), physiological stress (for example, starvation), or invading particles. These double-membrane-structured autophagosomes can be monitored via TEM. Alternatively, the increase of punctuated LC3-II is another sign of autophagosome accumulation. (2) Autophagosomes transfer to and fuse with lysosome to form autolysosome. (3) The wrapped contents are degraded in the autolysosome and the products (for example, amino acid) are released into the cytoplasm and recycled.

To understand the relationship between nanomaterials and autophagy, we need to monitor the autophagic activity accurately. There are some convenient methods to measure the number of autophagosomes including electron microscopy (EM) image analysis, fluorescent GFP-LC3 dots, or LC3 lipidation on a western blot. These methods are widely applied in current research of nanomaterial-induced autophagy. However, it is important to point out that autophagy is a dynamic, multi-step process that can be modulated at several steps. Accumulation of autophagosomes can reflect either an increase in autophagic activity or a blockage in the turnover of autophagosomes. Therefore, in addition to above discribed “static measurements”, some “autophagic

flux” assays are developed to distinguish these two possibilities, including LC3 turnover assay, GFP-LC3 cleavage assay, and degradation of long-lived proteins.

Autophagy is a tightly regulated process and increasing evidences demonstrate that autophagy may be involved in the progress of many diseases, including cancer, neurodegeneration, liver disease, muscular disorder and pathogen infection [27]. The relationship between autophagy and disease is often complicated. Autophagy can help cells to survive some stress conditions, such as nutrient and oxygen depletion [28,29]; however, autophagy can also promote autophagic cell death after chemotherapies. This is called “double-edged sword” effect of autophagy [27,30]. Therefore, it is crucial to take autophagy into consider in the study of biological effects of nanomaterials, since they may cause uncovered damage to organism through interfering with autophagic process [31]. On the other hand, it also implies that nanomaterials have the potential to be utilized in combination with chemotherapies for the treatment of diseases.

2 Carbon-based nanomaterials

Many forms of carbon-based nanomaterials were manufactured, including fullerenes C_{60} , carbon nanotubes (CNTs), and graphene sheet, carbon dots, nanodiamonds, nanonions and so on. Out of them, the former three types of nanomaterials are most well developed and widely used because of their unique properties in thermal stability, conductivity, mechanical and optical performance. Application of carbon-based nanomaterials is also a hot spot in biomedicine research [32–35]. There are a large number of reports about the cellular effect of carbon-based nanomaterials. Recently, several studies indicate that carbon-based nanomaterials can induce autophagy in different cell types (Table 1).

2.1 C_{60} and C_{60} derivatives

As early as in 2006, Yamawaki and Iwai [36] has found that the treatment of human umbilical vein endothelial cells (HUVECs) with $100 \mu\text{g/mL } C_{60}(\text{OH})_{24}$ for 24 h caused extensive internalization of fullerene and vacuolization. They observed aggregated fullerene in autophagosome via TEM. Protein immunoblot analysis of marker protein LC3 verified the occurrence of autophagy. The authors proposed that autophagy could be responsible at least partially for the cell death and injury caused by water-soluble fullerene, which represent a risk for atherosclerosis and ischemic heart disease. Similarly, Stephan et al. [40] showed that cell death upon exposure to $C_{60}(\text{OH})_{24}$ was related with cytoskeleton disruption, loss of mitochondrial capacity and autophagic vacuole accumulation. Indeed, ATP depletion and loss of mitochondrial potential were partially ameliorated when

Table 1 Autophagy induced by carbon-based nanomaterials

Nanomaterials	Average size (nm)	Cell lines	Autophagy marker	Reference
C ₆₀ (OH) ₂₄	7.1±2.4	HUVECs	TEM, LC3	[36]
Nano-C ₆₀	100	C6 cells	Acidophilic vesicles	[37]
Nano-C ₆₀	20–100	GFP-LC3/HeLa cells; Primary MEF cell	Acidic vesicles, LC3, Atg5	[38]
C ₆₀ (Nd) _x C ₆₀	50–100	HeLa cells; MCF-7; MEF cell	LC3, Atg5	[39]
C ₆₀ OH _x	20	LLC-PK1 cells	TEM, LC3	[40]
C ₆₀ -derived NPs	~78.9	Neuro-2A cells	LC3	[41]
SWNTs	~1	A549	TEM, LC3, Akt–TSC2–mTOR signaling	[42]
Graphene quantum dots	56.6±8.7	U251 cells	TEM, LC3, p62	[43]
Graphene oxide	2.4 μm and 350 nm	RAW264.7 cells	TEM, LC3	[44]

cells were co-treated with C₆₀(OH)₂₄ and an autophagy inhibitor (3-methyladenine).

On one hand, fullerene-induced autophagy can cause cell death or functional damage, leading to the development of certain diseases [36,40]. On the other hand, fullerene-triggered autophagy was found to play potential role in the treatment of diseases such as cancer and neurodegeneration. For example, Harhaji et al. [37] showed that the anti-glioma effect of low-dose nC₆₀ (0.25 μg/mL) was attenuated by pharmacological inhibition of autophagy. Hence, the authors proposed that autophagy might be involved in the anti-proliferative function of nC₆₀. Zhang et al. [38] systematically studied the effect of water-dispersed nanocrystal C₆₀ and its derivative C₆₀(Nd) nanoparticles on autophagy. Their results indicated that nC₆₀ could induce autophagy and sensitize chemotherapeutic killing of both normal and drug-resistant cancer cells at noncytotoxic concentrations (0.5 μg/mL). Nano-C₆₀-induced autophagy was reactive oxygen species (ROS)-dependent and enhanced by photo activation. The chemo-sensitization effect of nano-C₆₀ was autophagy-mediated and required functional Atg5, a key player in the autophagic signaling pathway. Compared to nC₆₀, nC₆₀(Nd) is more effective to induce autophagy and to sensitize chemotherapy [39]. These studies imply a novel application of nC₆₀/C₆₀(Nd) nanoparticle in cancer therapy.

Lee et al. [41] reported that fullerene-derived nanomaterials (PTX-C₆₀-2 and PEG-C₆₀-3) could elicit cytoprotective effect partially through autophagy to eliminate the accumulation of β-amyloid (Aβ) 25-35 in Neuro-2A cells. PTX-C₆₀-2 was more effective to induce autophagy compared to PEG-C₆₀-3. It is well known that aggregated β-amyloid is an important contributing factor in neuronal dysfunction and cellular apoptosis. Therefore the above study suggests PTX-C₆₀-2 has a potential in the treatment of neurodegenerative diseases.

2.2 Carbon nanotubes

Carbon nanotubes (CNTs) are mainly consisted of multi-walled carbon nanotubes (MWCNTs) and single-walled carbon nanotubes (SWCNTs). It is commonly accepted that

CNTs are toxic to cells and can lead to oxidative stress, inflammation, apoptosis and even necrosis [45–48]. Recently, autophagy was added to the list of CNTs-related cytotoxicity. Liu et al. [42] designed three types of functional single-walled carbon nanotubes (f-SWCNTs), including COOH-CNT, poly-aminobenzene sulfonic acid (PABS-CNT), PEG-CNT and studied the autophagic response and related pathway both *in vitro* and *in vivo*. They demonstrated that only COOH-CNT could trigger formation of autophagosomes and LC3-II up-regulation in A549 cells, indicating that functionalized surface can greatly affect the ability of nanomaterials to induce autophagy. Moreover, they found that COOH-CNT induce autophagic cell death in A549 cells through the AKT-TSC2-mTOR pathway and caused acute lung injury *in vivo*.

2.3 Graphene

Graphene and its oxidized forms (graphene oxide and reduced graphene oxide) refer to a single layer of carbon with a honeycomb structure and have sparked growing interest since 2004. Graphene has been applied in fields such as electrical, mechanical and biomedicine owing to its excellent physicochemical properties and well biocompatibility [49–51]. Therefore, more and more studies started focusing on the biosafety and cytotoxicity of graphene. Recently, Chen et al. [44] demonstrated that graphene oxide (GO) could induce autophagy in macrophage RAW264.7 cells and the effect was concentration-dependent. They observed the appearance of autophagic vacuoles and activation of LC3-II. Additionally, they uncovered that GO can simultaneously induce autophagy and TLR4/TLR9-regulated inflammatory, and the autophagic response was at least partly mediated by the TLR pathways.

Graphene quantum dots (GQD) possess both the unique physicochemical properties of graphene and the optical properties of quantum dots [52,53]. Markovic et al. [43] found that GQD irradiated with blue light (470 nm, 1 W) caused oxidative stress and kill U251 human glioma cells. Their results showed that the cell death triggered by photoexcited GQD was associated with both apoptosis and

autophagy. In addition to the accumulation autophagosomes, they further confirmed the up-regulation of autophagic flux by both the increase of LC3B-II and the markedly down-expression of p62, a protein selectively degraded by autophagy [54]. These data suggested potential usefulness of GQD in photodynamic therapy, but also increase more concerns about their possible toxicity.

3 Metal-based nanomaterials

Metal and their oxides have a long history to be applied in basically every aspects of life. With the development of nanotechnology, metal-based nanomaterials have been largely developed and manufactured, and are getting more and more commonly used in bioimaging, drug delivery, biosensors, and photo/thermal therapy. Undesired side effects of these metal-based nanoparticles are always hot topic in nano-related research. Certain metal-based nanomaterials like silver and copper oxide nanoparticles are known to be highly toxic due to oxidative damages. Oxidative stress is a common cause to autophagy [55,56]. Recently, several studies found that metal and metal oxide nanoparticles could cause autophagy *in vitro* and *in vivo* (Table 2).

3.1 Gold nanomaterials

Gold nanoparticles (AuNPs) are well known to be nearly non-cytotoxic and widely used as nanocarriers for many bioactive molecules [65,66]. Li et al. [57] found that gold nanoparticles (AuNPs) induce autophagy in human lung fibroblasts MRC-5 cells. They have observed the formation of autophagosomes, as well as upregulation of autophagy-related proteins (LC3, Atg7). Meanwhile, they found that AuNPs treatment could generate significant oxidative damage, as indicated by lipid hydroperoxides assay, expression levels of stress-response genes and proteins and anti-oxidants rescue. They hypothesized that AuNPs caused

oxidative stress in cells, and autophagy may play an important role in the cellular defense against oxidative toxicity.

In another study, Ma et al. [58] proposed that AuNP actually inhibited the autophagy process *via* block the fusion between autophagosome and lysosome. They have also observed the the accumulation of autophagosomes and the processing of the marker protein LC3-II when cells were incubated with AuNPs. However, they further demonstrated that the substrate protein p62 is also upregulated, suggesting the blockage of autophagy flux rather than induction of autophagy. Accordingly, they found that internalized AuNPs eventually entered lysosomes, caused alkalization of lysosome pH, and impaired lysosome degradation activity. This finding is important since it point out that nanoparticles can either promote or inhibit the process of autophagy, both resulted in the similar accumulation of autophagosomes and upregulation of LC3-II. Therefore it is important to analyze the autophagy process as a dynamic flux in addition to the classic status assays.

3.2 Iron-related nanomaterials

Wu et al. [59] examined the cytotoxicity of nanoparticles with an iron core and gold shell (Fe@AuO), and found Fe@AuO could cause irreversible loss of mitochondrial membrane-potential (MMP) in cancer cells, but only a transient MMP decrease in healthy cells. The MMP loss was correlated with up-regulation of mitochondria-mediated autophagy (Mitophagy). They also demonstrated that autophagy was the main reason of cytotoxicity. Similarly, Khan et al. [67] found that iron oxide nanoparticles selectively elicited autophagic cell death in cancer cells (A549) but not in normal cells (IMR-90). In this case, autophagy was related with ROS production as well as mitochondrial damage. Their further study suggested that iron oxide-triggered autophagy is partially depended on the classical mTOR pathway. These results imply that Fe@AuO and iron oxide nanoparticles have the potential as therapeutic reagent to specifically induce autophagic cell death in tumor cells.

Table 2 Autophagy induced by metal-based nanomaterials

Nanomaterials	Average size (nm)	Cell lines	Autophagy marker	Reference
AuNPs	20	MRC-5 cells	TEM, LC3, Atg5, Atg 7, Atg 12	[57]
AuNPs	10, 25 and 50	NRK cells	TEM, LC3, p62	[58]
Fe@Au NPs	10	hNOK cells; OECM1 cells	TEM, LC3	[59]
Iron oxide NPs	51.34±14.71	A549 cells; IMR-90 cells	LC3, mTOR	[31]
Superparamagnetic iron oxide	8	HCECs cells	TEM, LC3	[60]
TiO ₂ NPs	<25	H4-LC3-GFP cells	LC3	[61]
TiO ₂ NPs	21	HCECs cells	TEM, LC3	[60]
α-Al ₂ O ₃ NPs	60; 200	Dendritic cells (DCs)	TEM, LC3, Atg5-Atg12	[62]
Nanoalumina	8–12	HCMECs/D3 cells; C57BL/6 mice	LC3, p62	[63]
MnNPs	25	N27 dopaminergic neuronal cells	TEM, LC3	[64]
AgNW	3 μm (length)	THP-1 cells	TEM, LC3	[24]

3.3 Alumina and its oxide nanoparticles

It has been reported that aluminum can cause neurodegenerative diseases, which are resulted from the presence of oxidative stress, inflammatory and the breakdown of the blood-brain barrier (BBB) [63,68,69]. Chen et al. [63] studied the toxicity and related mechanism of nanoalumina at both cellular and animal levels. They found that nanoalumina could pass through BBB, accumulate in the mice brain and locate into mitochondria in brain cells. Nanoalumina-induced dysfunction of mitochondria elicited autophagy, decreased tight-junction protein expression, and elevated BBB permeability. Therefore, autophagy may be a primary mechanism involved in nanoalumina-induced neurovascular toxicity in the central nervous system.

In another study, Li et al. [62] showed that α -Al₂O₃ nanoparticles at two sizes (60 and 200 nm) could both induce autophagy effectively and exhibited potent antitumor capability. α -Al₂O₃ nanoparticles function as an antigen carrier to T cells through the autophagy pathway, which decreased the needed amount of antigen to stimulate production of enough T-cells. Administration of α -Al₂O₃ NPs incubated with the autophagic vacuoles collected from the tumor cells greatly enhanced the effect of cancer therapy.

3.4 Others metal-based nanomaterials

Titanium dioxide (TiO₂) is widely applied as a supplement in cosmetics, coating and paint due to its high refractive index and bold white coloration [61]. Therefore, the cytotoxicity of titanium dioxide attracted much attention. In a parallel study including different metal oxide NPs like ZnO, FeO and TiO₂, Yu et al. [61] found that only TiO₂ can effectively induce autophagy, as measured by the activation of LC3 and the increase of autophagic flux. Similarly, Blanka et al. [60] demonstrated that TiO₂-induced autophagy was mainly related with NPs aggregation and oxidative stress. They also found that nanomaterials-mediated autophagy was size-dependent.

In addition, it has been reported that silver nanowire (AgNW), manganese nanoparticles (MnNPs) could effectively elicit autophagy in epithelial, endothelial, gastric, phagocytic cells [24] and dopaminergic neuronal cells [64], respectively.

4 Rare earth oxides nanomaterials

Rare earth elements refer to a set of seventeen chemical elements including fifteen lanthanides plus scandium and yttrium in the periodic table. Rare earth oxides naturally exist in the earth crust. Owing to their unique chemical and physical properties, rare earth oxides materials are widely applied in metallurgy, ceramics, magnets laser industries and biological field [70]. With the growing application, the biosafety issue of rare earth oxides materials has caused great concerns, especially in the field of nanoscale research.

Wen group [71–74] in University of Science and Technology of China made a series of excellent study on the toxicity of rare earth oxides nanomaterials, mainly focused on their impact on cellular autophagy (Table 3). They found that micromolar equivalent concentration of nano-sized neodymium oxide (Nano Nd₂O₃) could induce autophagy and cell death in non-small cell lung cancer NCI-H460 cells. They have observed the massive vacuolization, S-phase cell cycle arrest, loss of mitochondrial membrane potential (MMP) and inhibition of proteasome activity, suggesting that cell death was resulted from autophagy [71]. Subsequently, they demonstrated that all of four different kinds of rare earth oxides nanomaterials, including samarium oxide (Nano Sm₂O₃), europium oxide (Nano Eu₂O₃), gadolinium oxide (Nano Gd₂O₃) and terbium oxide (Nano Tb₂O₃), could induce accumulation of GFP-LC3 punctate in HeLa cells stable expressing GFP-LC3 fusion proteins. And the autophagic response was dose- and time-dependent [72]. In another study, they investigated three rare earth oxides nanomaterials, Yttrium oxide (Y₂O₃), Ytterbium (Yb₂O₃) and Lanthanum (La₂O₃). Their results demonstrated that all of

Table 3 Autophagy induced by rare earth oxides nanomaterials

Nanomaterials	Average size (nm)	Cell lines	Autophagy marker	Reference
Nano Nd ₂ O ₃	80	Human NCI-H460 cell line	Vacuoles	[71]
Nano Eu ₂ O ₃	~50	GFP-LC3/HeLa cells	TEM, LC3	[72]
Nano Sm ₂ O ₃	~20	GFP-LC3/HeLa cells	TEM, LC3	[72]
Nano Gd ₂ O ₃	~100	GFP-LC3/HeLa cells	TEM, LC3	[72]
Nano Tb ₂ O ₃	~80	GFP-LC3/HeLa cells	TEM, LC3	[72]
Y ₂ O ₃	<50	GFP-LC3/HeLa cells Atg5 ^{-/-} MEF cells Normal MEF cells	Vacuoles, LC3, p62, Atg5	[73]
Yb ₂ O ₃	<50	GFP-LC3/HeLa cells Atg5 ^{-/-} MEF cells Normal MEF cells	Vacuoles, LC3, p62, Atg5	[73]
LN-based nanocrystals	~20	GFP-LC3/HeLa cells	TEM, LC3	[74]
CeO ₂ NPs	10–30	CD14 ⁺ cells	TEM, LC3B	[15]

them could trigger autophagy but only Y_2O_3 and Yb_2O_3 cause vacuolization, suggesting that commonly observed vacuolization differs from autophagic pathway [73]. More recently, Zhang et al. [74] achieved to control the ability of Ln_2O_3 nanocrystals to induce autophagy via surface-coating peptides. In a screening assay they identified a short peptide RE-1 that could reduce Ln_2O_3 induced autophagy. Furthermore, adding an RGD motif to RE-1 could effectively target the nanocrystals to tumors and increase autophagic cell death *in vivo*. This work discovered a powerful tool to regulate the nanomaterial-cell interactions and to achieve the desired level of autophagy, which may promote the application of rare earth oxides nanomaterials in cancer diagnose and therapy.

Besides, Hussain et al. [75] reported cerium dioxide (CeO_2) nanoparticles could simultaneously induce apoptosis and autophagy, resulting in cell death of human monocytes. The autophagy effect induced by CeO_2 is time- and dose-dependent, and modulated by p53. However, the detailed mechanism is unclear.

5 Semiconductor quantum dots

Quantum dots (QDs) are semiconductor nanocrystals with a typical diameter ranging from 2 to 10 nm. QDs possess superior optical properties, such as high photoluminescence quantum yield (PLQY), broad absorption coupled with narrow emission, and strong photostability [76]. Consequently, several kinds of QDs have been fabricated and utilized for applications including biosensing, bioimaging and disease diagnosis [77]. However, Most QDs are made of heavy metal ions, which may result in potential toxicity that hampers their practical applications. And some heavy metal ions can induce autophagic cell death. Therefore, in the study of QDs-related cellular effects, it is necessary to inspect the change of autophagy levels [78].

Seleverstov et al. [79] studied the process of internalization and compared the cytotoxicity of two different-sized QDs (605 and 525 nm) in the human mesenchymal stem

cells (hMSC). In the cells treated with QDs of smaller size (QD525), they observed autophagic cell death and the massive formation of autophagosomes containing damaged organelles and aggregated QDs after 72 h. In contrast, little autophagic vacuoles or damaged mitochondria were found in hMSC cells labeled with QDs of bigger size (QD605). These results for the first time suggested that the ability of nanoparticles to induce autophagy is size-dependent.

In another study, Stern et al. [78] compared the cytotoxicity on LLC-PK1 porcine kidney cells of two QDs species, which have similar size but are made of different core materials (CdSe and InGaP). The core size of CdSe and InGaP was 5.1 and 3.7 nm, respectively. Both QDs can induce autophagy as demonstrated by LC3 immunoblotting, TEM pictures and LysoTracker staining, suggesting that autophagy is not caused directly by certain metal element in the composition. Besides, they also found that CdSe QD elicited stronger autophagic response than InGaP QD at equal molar concentrations, which is positively correlated to their cytotoxicity [78].

6 Summary and outlook

Autophagy was originally defined as type II programmed cell death. With the detailed study, it is more and more clear that autophagy plays fundamental roles in multiple cellular processes and is closely associated with cancer and neurodegenerative disease. Particularly, rapamycin, a classic drug in the treatment of cancer and metabolic disease, is the central regulator of autophagy. Therefore, manipulation of autophagy levels with nanomaterials provides novel potential opportunities for cancer therapy.

Lately, a variety of nanomaterials have been reported as autophagy inducers. However, in many studies, the accumulation of autophagosome protein LC3II was commonly used as a marker of up-regulated autophagy. However, the process of autophagy is a dynamic flux (Figure 1). It is essential to carefully examine and distinguish between the "upstream induction" of autophagosome formation and the




Figure 1 Nanomaterials interfere with the process of autophagy.

“downstream blockage” of fusion between autophagosome and lysosome, which result in similar phenotype but completely different biological consequence. Gold nanoparticles, for example, were firstly reported to induce autophagy, which turned out to be inhibitor of the acidification of lysosome and turnover of autophagosomes by “autophagic flux” assays. Such incompleteness also exist in current study of other nanomaterials and autophagic activity. Therefore it is essential to perform multiple assays to verify an autophagic response. In future studies, we suggest performing studies to better understand detailed interactions between nanomaterials and autophagic pathways and to uncover the underlying mechanism.

This work was supported by the National Basic Research Program of China (2011CB943900) and the National Natural Science Foundation of China (31071145).

- Lundqvist M, Stigler J, Elia G, et al. Nanoparticle size and surface properties determine the protein corona with possible implications for biological impacts. *P Natl Acad Sci USA*, 2008, 105: 14265–14270
- Chomoucka J, Drbohlavova J, Huska D, et al. Magnetic nanoparticles and targeted drug delivering. *Pharmacol Res*, 2010, 62: 144–149
- Häfeli U O, Riffle J S, Harris-Shekhawat L, et al. Cell uptake and *in vitro* toxicity of magnetic nanoparticles suitable for drug delivery. *Mol Pharmaceut*, 2009, 6: 1417–1428
- Petrenko V A. Evolution of phage display: From bioactive peptides to bioselective nanomaterials. *Expert Opin Drug Del*, 2008, 5: 825–836
- Aitken R, Chaudhry M, Boxall A, et al. Manufacture and use of nanomaterials: Current status in the UK and global trends. *Occup Med-Oxford*, 2006, 56: 300–306
- Helland A, Wick P, Koehler A, et al. Reviewing the environmental and human health knowledge base of carbon nanotubes. *Environ Health Persp*, 2007, 15: 1125–1131
- Marambio-Jones C, Hoek E M. A review of the antibacterial effects of silver nanomaterials and potential implications for human health and the environment. *J Nanopart Res*, 2010, 12: 1531–1551
- Thomas K, Sayre P. Research strategies for safety evaluation of nanomaterials, Part I: Evaluating the human health implications of exposure to nanoscale materials. *J Nanopart Res*, 2005, 87: 316–321
- Xia T, Kovoichich M, Brant J, et al. Comparison of the abilities of ambient and manufactured nanoparticles to induce cellular toxicity according to an oxidative stress paradigm. *Nano Lett*, 2006, 6: 1794–1807
- Li N, Venkatesan M I, Miguel A, et al. Induction of heme oxygenase-1 expression in macrophages by diesel exhaust particle chemicals and quinones via the antioxidant-responsive element. *J Immunol*, 2000, 165: 3393–3401
- Braydich-Stolle L K, Schaeublin N M, Murdock R C, et al. Crystal structure mediates mode of cell death in TiO₂ nanotoxicity. *J Nanopart Res*, 2009, 11: 1361–1374
- Donaldson K, Stone V, Tran C, et al. Nanotoxicology. *Occup Environ Med*, 2004, 61: 727–728
- Stern S T, Johnson D N. Role for nanomaterial-autophagy interaction in neurodegenerative disease. *Autophagy*, 2008, 4: 1097–1100
- Stern S, Adisheshaiah P, Crist R. Autophagy and lysosomal dysfunction as emerging mechanisms of nanomaterial toxicity. *Part Fibre Toxicol*, 2012, 9: 20
- Hussain S, Al-Nsour F, Rice A B, et al. Cerium dioxide nanoparticles induce apoptosis and autophagy in human peripheral blood monocytes. *ACS Nano*, 2012, 6: 5820–5829
- Sohaebuddin S K, Thevenot P T, Baker D, et al. Nanomaterial cytotoxicity is composition, size, and cell type dependent. *Part Fibre Toxicol*, 2010, 7: 22
- Zhang Q, Yang W, Man N, et al. Autophagy-mediated chemosensitization in cancer cells by fullerene C₆₀ nanocrystal. *Autophagy*, 2009, 5: 1107–1117
- He C, Klionsky D J. Regulation mechanisms and signaling pathways of autophagy. *Annu Rev Genet*, 2009, 43: 67
- Mizushima N. Autophagy in protein and organelle turnover. *Cold Spring Harb Symp Quant Biol*, 2011, 76: 397–402
- De Stefano D, Carnuccio R, Maiuri M C. Nanomaterials toxicity and cell death modalities. *J Drug Deliv*, 2012, 2012: 1–12
- Mizushima N, Yoshimori T, Levine B. Methods in mammalian autophagy research. *Cell*, 2010, 140: 313–326
- Zabirnyk O, Yezhelyev M, Seleverstov O. Nanoparticles as a novel class of autophagy activators. *Autophagy*, 2007, 3: 278–281
- Li C, Liu H, Sun Y, et al. PAMAM nanoparticles promote acute lung injury by inducing autophagic cell death through the Akt-TSC2-mTOR signaling pathway. *J Mol Cell Biol*, 2009, 1: 37–45
- Verma N K, Conroy J, Lyons P E, et al. Autophagy induction by silver nanowires: A new aspect in the biocompatibility assessment of nanocomposite thin films. *Toxicol Appl Pharm*, 2012, 264: 451–461
- Sun T T, Yan Y W, Zhao Y, et al. Copper oxide nanoparticles induce autophagic cell death in A549 cells. *PLoS One*, 2012, 7: 1–7
- De Duve C. The lysosome. *Sci Am*, 1963, 208: 64
- Shintani T, Klionsky D J. Autophagy in health and disease: A double-edged sword. *Science*, 2004, 306: 990–995
- Rabinowitz J D, White E. Autophagy and metabolism. *Science*, 2010, 330: 1344–1348
- Degenhardt K, Mathew R, Beaudoin B, et al. Autophagy promotes tumor cell survival and restricts necrosis, inflammation, and tumorigenesis. *Cancer Cell*, 2006, 10: 51–64
- Yu L, Strandberg L, Lenardo M J. The selectivity of autophagy and its role in cell death and survival. *Autophagy*, 2008, 4: 567–573
- Khan M I, Mohammad A, Patil G, et al. Induction of ROS, mitochondrial damage and autophagy in lung epithelial cancer cells by iron oxide nanoparticles. *Biomaterials*, 2012, 33: 1477–1488
- Bianco A, Kostarelos K, Prato M. Applications of carbon nanotubes in drug delivery. *Curr Opin Chem Biol*, 2005, 9: 674
- Lin Y, Taylor S, Li H, et al. Advances toward bioapplications of carbon nanotubes. *J Mater Chem*, 2004, 14: 527–541
- Wong S, Kam N, Dai H. Single walled carbon nanotubes for transport and delivery of biological cargos. *Phys Status Solidi B*, 2006, 243: 3561–3566
- Bianco A, Kostarelos K, Prato M. Opportunities and challenges of carbon-based nanomaterials for cancer therapy. *Expert Opin Drug Del*, 2008, 5: 331–342
- Yamawaki H, Iwai N. Cytotoxicity of water-soluble fullerene in vascular endothelial cells. *Am J Physiol-Cell Ph*, 2006, 290: C1495–C1502
- Harhaji L, Isakovic A, Raicevic N, et al. Multiple mechanisms underlying the anticancer action of nanocrystalline fullerene. *Eur J Pharmacol*, 2007, 568: 89–98
- Zhang Q, Yang W, Man N, et al. Autophagy-mediated chemosensitization in cancer cells by fullerene C₆₀ nanocrystal. *Autophagy*, 2009, 5: 1107–1117
- Wei P F, Zhang L, Lu Y, et al. C₆₀(Nd) nanoparticles enhance chemotherapeutic susceptibility of cancer cells by modulation of autophagy. *Nanotechnology*, 2010, 21: 1–12
- Johnson-Lyles D N, Peifley K, Lockett S, et al. Fullerenol cytotoxicity in kidney cells is associated with cytoskeleton disruption, autophagic vacuole accumulation, and mitochondrial dysfunction. *Toxicol Appl Pharm*, 2010, 248: 249–258
- Lee C M, Huang S T, Huang S H, et al. C₆₀ fullerene-pentoxifylline dyad nanoparticles enhance autophagy to avoid cytotoxic effects caused by the β -amyloid peptide. *Nanomedicine*, 2011, 7: 107–114
- Liu H L, Zhang Y L, Yang N, et al. A functionalized single-walled carbon nanotube-induced autophagic cell death in human lung cells through Akt-TSC2-mTOR signaling. *Cell Death Dis*, 2011, 2: 1–7
- Markovic Z M, Ristic B Z, Arskin K M, et al. Graphene quantum dots as autophagy-inducing photodynamic agents. *Biomaterials*, 2012, 33: 7084–7092

- 44 Chen G Y, Yang H J, Lu C H, et al. Simultaneous induction of autophagy and toll-like receptor signaling pathways by graphene oxide. *Biomaterials*, 2012, 33: 6559–6569
- 45 Sharma C S, Sarkar S, Periyakaruppan A, et al. Single-walled carbon nanotubes induces oxidative stress in rat lung epithelial cells. *J Nanosci Nanotechnol*, 2007, 7: 2466–2472
- 46 Panessa-Warren B, Warren J, Wong S, et al. Biological cellular response to carbon nanoparticle toxicity. *J Phys-Condens Mat*, 2006, 18: S2185–S2201
- 47 Bottini M, Bruckner S, Nika K, et al. Multi-walled carbon nanotubes induce T lymphocyte apoptosis. *Toxicol Lett*, 2006, 160: 121–126
- 48 Lam C W, James J T, McCluskey R, et al. Pulmonary toxicity of single-wall carbon nanotubes in mice 7 and 90 days after intratracheal instillation. *J Nanopart Res*, 2004, 7: 126–134
- 49 Feng L, Liu Z. Graphene in biomedicine: Opportunities and challenges. *Nanomedicine*, 2011, 6: 317–324
- 50 Chen H, Müller M B, Gilmore K J, et al. Mechanically strong, electrically conductive, and biocompatible graphene paper. *Adv Mater*, 2008, 20: 3557–3561
- 51 Wu Z S, Ren W, Gao L, et al. Synthesis of graphene sheets with high electrical conductivity and good thermal stability by hydrogen arc discharge exfoliation. *ACS Nano*, 2009, 3: 411–417
- 52 Christensen I L, Sun Y P, Juzenas P. Carbon dots as antioxidants and prooxidants. *J Biomed Nanotechnol*, 2011, 7: 667–676
- 53 Agostinis P, Berg K, Cengel K A, et al. Photodynamic therapy of cancer: An update. *Ca-Cancer J Clin*, 2011, 61: 250–281
- 54 Ichimura Y, Kumanomidou T, Sou Y, et al. Structural basis for sorting mechanism of p62 in selective autophagy. *J Biol Chem*, 2008, 283: 22847–22857
- 55 AshaRani P V, Mun G L K, Hande M P, et al. Cytotoxicity and genotoxicity of silver nanoparticles in human cells. *ACS Nano*, 2008, 3: 279–290
- 56 Navarro E, Piccapietra F, Wagner B, et al. Toxicity of silver nanoparticles to *Chlamydomonas reinhardtii*. *Environ Sci Technol*, 2008, 42: 8959–8964
- 57 Li J J, Hartono D, Ong C N, et al. Autophagy and oxidative stress associated with gold nanoparticles. *Biomaterials*, 2010, 31: 5996–6003
- 58 Ma X, Wu Y, Jin S, et al. Gold nanoparticles induce autophagosome accumulation through size-dependent nanoparticle uptake and lysosome impairment. *ACS Nano*, 2011, 5: 8629–8639
- 59 Wu Y N, Yang L X, Shi X Y, et al. The selective growth inhibition of oral cancer by iron core-gold shell nanoparticles through mitochondria-mediated autophagy. *Biomaterials*, 2011, 32: 4565–4573
- 60 Blanka H K, Catherine C B, Seher G A, et al. Induction of oxidative stress, lysosome activation and autophagy by nanoparticles in human brain-derived endothelial cells. *Biochem J*, 2012, 441: 813–821
- 61 Julia X Y, Li T H, Yu J, et al. Distinct biological effects of different nanoparticles commonly used in cosmetics and medicine coatings. *Cell Biosci*, 2011, 1: 19
- 62 Li H, Li Y, Jiao J, et al. Alpha-alumina nanoparticles induce efficient autophagy-dependent cross-presentation and potent antitumour response. *Nat Nanotechnol*, 2011, 6: 645–650
- 63 Chen L, Zhang B, Toborek M. Autophagy is involved in nanoalumina-induced cerebrovascular toxicity. *Nanomed-Nanotechnol*, 2013, 9: 212–221
- 64 Afeseh N H, Kanthasamy A, Gu Y, et al. Manganese nanoparticle activates mitochondrial dependent apoptotic signaling and autophagy in dopaminergic neuronal cells. *Toxicol Appl Pharm*, 2011, 256: 227–240
- 65 Wei M, Chen N, Li J, et al. Polyvalent immunostimulatory nanoagents with self-assembled CpG oligonucleotide-conjugated gold nanoparticles. *Angew Chem Int Ed Engl*, 2012, 51: 1202–1206
- 66 Wang B, Chen N, Wei Y, et al. Akt signaling-associated metabolic effects of dietary gold nanoparticles in *Drosophila*. *Sci Rep*, 2012, 2: 563
- 67 Khan M I, Mohammad A, Patil G, et al. Induction of ROS, mitochondrial damage and autophagy in lung epithelial cancer cells by iron oxide nanoparticles. *Biomaterials*, 2012, 33: 1477–1488
- 68 Bondy S C. The neurotoxicity of environmental aluminum is still an issue. *Neurotoxicology*, 2010, 31: 575–581
- 69 Oesterling E, Chopra N, Gavalas V, et al. Alumina nanoparticles induce expression of endothelial cell adhesion molecules. *Toxicol Lett*, 2008, 178: 160–166
- 70 Man N, Yu S H. Rare earth oxide nanocrystals as a new class of autophagy inducers. *Autophagy*, 2010, 6: 310–311
- 71 Chen Y, Yang L, Feng C, et al. Nano neodymium oxide induces massive vacuolization and autophagic cell death in non-small cell lung cancer NCI-H460 cells. *Biochem Biophys Res Commun*, 2005, 337: 52–60
- 72 Yu L, Lu Y, Man N, et al. Rare earth oxide nanocrystals induce autophagy in HeLa cells. *Small*, 2009, 5: 2784–2787
- 73 Zhang Y, Yu C, Huang G, et al. Nano rare-earth oxides induced size-dependent vacuolization: An independent pathway from autophagy. *Int J Nanomedicine*, 2010, 5: 601–609
- 74 Zhang Y, Zheng F, Yang T, et al. Tuning the autophagy-inducing activity of lanthanide-based nanocrystals through specific surface-coating peptides. *Nat Mater*, 2012, 11: 817–826
- 75 Hussain S, Garantzotis S. Interplay between apoptotic and autophagy pathways after exposure to cerium dioxide nanoparticles in human monocytes. *Autophagy*, 2013, 9: 101–103
- 76 He Y, Lu H T, Sai L M, et al. Microwave synthesis of water-Dispersed CdTe/CdS/ZnS core-shell-Shell quantum dots with excellent photostability and biocompatibility. *Adv Mater*, 2008, 20: 3416–3421
- 77 Peng C W, Li Y. Application of quantum dots-based biotechnology in cancer diagnosis: Current status and future perspectives. *J Nanomater*, 2010, 2010: 1–11
- 78 Stern S T, Zolnik B S, McLeland C B, et al. Induction of autophagy in porcine kidney cells by quantum dots: A common cellular response to nanomaterials? *J Nanopart Res*, 2008, 10: 140–152
- 79 Seleverstov O, Zabirnyk O, Zscharnack M, et al. Quantum dots for human mesenchymal stem cells labeling. A size-dependent autophagy activation. *Nano Lett*, 2006, 6: 2826–2832

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.